

# INTRAVENOUS CANCER TREATMENT EDUCATION



PASSION FOR PATIENTS  
NCODA.ORG



## MOGAMULIZUMAB

### Name of the regimen and cancer drugs

Mogamulizumab-kpkc (moh GAM yoo LIH zoo mab): Poteligeo®

### Common uses

This regimen is most commonly used for mycosis fungoides or Sezary syndrome but may be used for other diagnoses.

### Treatment schedule

Your treatment will be given into your vein through an intravenous (IV) line. This may be into a short, flexible temporary catheter in your arm, or through a central venous catheter. A central venous catheter, or central line is a long, flexible IV tube that empties into a very large vein next to the heart. Talk with your care team to see which will be best for you and your treatment.

Each mogamulizumab treatment is repeated every 28 days. This is known as one cycle. Your treatment may be given for a set number of cycles, or it will keep going until the drug, or drugs, stop working or you have side effects which stop you from continuing treatment.

#### First Cycle

- Mogamulizumab IV given on days 1, 8, 15, and 22

Drug	Cycle 1 Day	1	...	8	...	15	...	22	...	28	Cycle 2 Day 1
Mogamulizumab											

#### Cycle 2 and Beyond

- Mogamulizumab IV given on days 1 and 15

Drug	Cycle 1 Day	1	...	8	...	15	...	22	...	28	Cycle 2 Day 1
Mogamulizumab											

### Other medications

Other medications may be ordered for you to prevent or treat certain side effects. These include:

	Instructions:
<b>Medications to prevent allergic reactions</b>	<p>You will receive medications to prevent infusion related reactions prior to starting your cancer treatment.</p> <p>_____</p> <p>_____</p> <p>_____</p> <p>_____</p>

# INTRAVENOUS CANCER TREATMENT EDUCATION



## MOGAMULIZUMAB

### Possible drug interactions

- Mogamulizumab may interact with other drugs you are taking. Please inform your care providers of all prescription medicines, over-the-counter medications, vitamins, and herbal products that you take.
- Talk with your care provider or pharmacist before taking new medications, supplements, or receiving any vaccines.

### Common Side Effects

Common side effects that have been known to happen in patients receiving mogamulizumab are listed in the left side of this table. In some instances, the side effects may be reported less often but are still important to discuss. This table does not list all the known side effects for this therapy, only the ones that are experienced most often. Not every patient experiences every known side effect of a drug; even if you are taking the same drug as another patient, you may experience different side effects. Options to help manage any side effects that do occur are included on the right side of this table. These should be discussed with your care provider. If you experience any side effect you cannot manage or that is not listed here, contact your care provider.

Possible Side Effect	Management
<p><b>Decreased hemoglobin, part of the red blood cells that carry iron and oxygen</b></p>	<p>Your hemoglobin should be checked by a simple blood test. When your hemoglobin is low, you may notice that you get tired or fatigued more easily.</p> <ul style="list-style-type: none"> <li>• Try to get 7 to 8 hours of sleep per night</li> <li>• Avoid operating heavy machinery if you feel too tired</li> <li>• Find a balance between “work” and “rest”</li> <li>• Stay as active as possible, but know that it is okay to rest as needed, too</li> <li>• You might notice that you are more pale than usual</li> </ul> <p>Let your care team know right away if you have:</p> <ul style="list-style-type: none"> <li>• Shortness of breath</li> <li>• Dizziness</li> <li>• Fast or abnormal heartbeat</li> <li>• Headache</li> </ul>
<p><b>Decreased white blood cells (WBCs) and increased risk for infection</b></p>	<p>Your WBCs should be monitored by a simple blood test. When your WBCs are low, you are at a greater risk of having an infection. Take the following precautions to protect yourself from infection:</p> <ul style="list-style-type: none"> <li>• Wash your hands often, especially before eating and after using the bathroom.</li> <li>• Avoid crowds and people with fevers, flu, or other infection.</li> <li>• Bathe often for good personal hygiene.</li> </ul> <p>Contact your care team if you experience any signs or symptoms of an infection such as:</p> <ul style="list-style-type: none"> <li>• Fever (temperature more than 100.4°F or 38°C)</li> <li>• Chills</li> <li>• Sore throat</li> <li>• Burning when peeing</li> <li>• Tiredness that is worse than normal</li> <li>• A sore that becomes red, is draining, or does not heal</li> </ul> <p>Check with your care team before taking any medicine for a fever or chills.</p>

*Continued on the next page*

# INTRAVENOUS CANCER TREATMENT EDUCATION



## MOGAMULIZUMAB

Possible Side Effect	Management
<p><b>Decreased platelet count and increased risk of bleeding</b></p>	<p>Your platelets should be monitored by a simple blood test. When they are low, you may bruise or bleed more easily than usual.</p> <ul style="list-style-type: none"> <li>• Use caution to avoid bruises, cuts, or burns</li> <li>• Blow your nose gently and do not pick your nose</li> <li>• Brush your teeth gently with a soft toothbrush and maintain good oral hygiene</li> <li>• When shaving use an electric razor instead of razor blades</li> <li>• Use a nail file instead of a nail clippers</li> </ul> <p>Call your care team if you have bleeding that won't stop. Examples include:</p> <ul style="list-style-type: none"> <li>• A bloody nose that bleeds for more than 5 minutes despite pressure</li> <li>• A cut that continues to ooze despite pressure</li> <li>• Gums that bleed a lot when you floss or brush</li> </ul> <p>Seek medical help right away if you have any severe headaches, blood in your urine or stool, coughing up blood, or bleeding that you cannot stop or lasts a long time.</p> <p><b>You may need to take a break or “hold” your medication for medical or dental procedures. Talk to your care team or dentist before any scheduled procedures.</b></p>
<p><b>Changes in electrolytes and other laboratory values</b></p> <ul style="list-style-type: none"> <li>• High glucose levels</li> <li>• Decreased albumin levels</li> <li>• Decreased calcium levels</li> <li>• Increased uric acid</li> </ul>	<p>Changes in some lab values may occur and will be monitored by a simple blood test.</p> <ul style="list-style-type: none"> <li>• You may not feel any symptoms if the changes are mild and they usually are not a sign of a serious problem</li> <li>• More severe changes may occur which can be a sign of a serious problem</li> </ul> <p>Notify your care team if you have any of the following:</p> <ul style="list-style-type: none"> <li>• Shortness of breath</li> <li>• Chest discomfort</li> <li>• Weakness or fatigue</li> <li>• New aches and pains</li> <li>• Headaches</li> <li>• Dizziness</li> <li>• Swelling of your legs or feet</li> <li>• Red or brown colored urine</li> </ul>
<p><b>Rash or itchy skin</b></p>	<ul style="list-style-type: none"> <li>• Keep your skin moisturized with creams and moisturizing lotions to decrease the risk of rash or itchiness and wear loose fitting clothing.</li> <li>• Avoid using perfumes and cologne as these products may increase rash symptoms.</li> <li>• Avoid being in the heat for long periods of time.</li> <li>• Your provider may recommend an over-the-counter antihistamine or a topical cream.</li> <li>• Sunlight can make symptoms worse.             <ul style="list-style-type: none"> <li>• Avoid sun exposure as much as possible to decrease the risk of sunburn. The highest exposure to UV (ultra-violet) radiation occurs between the hours of 10am and 4pm.</li> </ul> </li> </ul>

*Continued on the next page*

# INTRAVENOUS CANCER TREATMENT EDUCATION



PASSION FOR PATIENTS  
NCODA.ORG



## MOGAMULIZUMAB

Possible Side Effect	Management
<p><b>Rash or itchy skin (continued)</b></p>	<ul style="list-style-type: none"> <li>• Sunlight can make symptoms worse.                             <ul style="list-style-type: none"> <li>• Wear long-sleeved clothing, with UV protection if possible.</li> <li>• Wear broad-brimmed hats.</li> <li>• Apply broad-spectrum sunscreen (UVA/UVB) with at least SPF 30 as often as directed on the bottle.</li> <li>• Use lip balm with at least SPF 30.</li> <li>• Avoid tanning beds.</li> </ul> </li> </ul> <p>If your rash or itching continues to worsen, contact your care team.</p>
<p><b>Infusion reactions</b></p>	<p>During your treatment, let the nurse know right away if any of these symptoms happen:</p> <ul style="list-style-type: none"> <li>• Chills or shaking</li> <li>• Dizziness</li> <li>• Fever</li> <li>• Itchiness or rash</li> <li>• Flushing</li> <li>• Difficulty breathing</li> <li>• Wheezing</li> <li>• Throat irritation</li> <li>• Cough</li> <li>• Congestion</li> <li>• Sudden back pain</li> <li>• Feeling faint</li> </ul>
<p><b>Fatigue</b></p>	<ul style="list-style-type: none"> <li>• You may be more tired than usual or have less energy</li> <li>• Stay as active as possible, but know it is okay to rest as needed</li> <li>• Try to do some type of moderate activity every day</li> <li>• Conserve your energy. Plan your activities and do them at a time of day when you feel a bit more energetic</li> <li>• Follow a healthy diet and stay hydrated</li> <li>• Accept help from family and friends</li> <li>• Find healthy ways to manage stress, such as meditation, journaling, yoga, and guided imagery</li> <li>• Develop good sleeping habits, limit napping during the day to help you sleep better at night</li> <li>• Avoid operating heavy machinery if you feel too tired</li> </ul> <p>Contact your care team if you experience extreme fatigue that prevents you from doing your normal daily activities</p>

# INTRAVENOUS CANCER TREATMENT EDUCATION



PASSION FOR PATIENTS  
NCODA.ORG



## MOGAMULIZUMAB

### Rare but serious side effects

Mogamulizumab rarely causes life-threatening and fatal immune-mediated reactions. Let your team know if you experience muscle pain or weakness, fatigue, swelling, difficulty breathing, dark-colored urine, or yellowing of the skin and eyes.

- Patients who receive an allogeneic stem cell transplant after mogamulizumab may experience serious transplant-related complications, which may even lead to death. Your healthcare provider will monitor you for signs of complications if you have a stem cell transplant. If you experience ANY new, worsening, or uncontrolled side effects, call your care team immediately.

If you experience ANY new, worsening, or uncontrolled side effects, call your care team immediately.

**(INSTITUTIONAL CONTACT INFO)**

### Handling body fluids and waste

Some of the drugs you receive may be present in urine, stool, sweat, or vomit for many days after treatment. Many cancer drugs are toxic, your bodily waste may also be toxic and can be dangerous to come into contact with. Once you have started mogamulizumab, follow the instructions below for at least two days after your treatment. This is to keep you, your loved ones, and the environment as safe as possible.

- Pregnant women should avoid touching anything that may be soiled with body fluids from the patient.
- Toilet and septic systems
  - You may use the same toilet, septic tank, and/or sewer that you usually use. If you have a low-flow toilet, close the lid and flush twice to ensure that all waste has been discarded.
- If the toilet or toilet seat becomes soiled with urine, stool, or vomit, clean the surface after every use before other people use the toilet.
- Wash hands with soap and water after using the toilet for at least 20 seconds.
- If you need a bedpan, be sure your caregiver knows to wear gloves to assist with cleanup and to wash the bedpan with soap and water every day.
- If you do not have good control of bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb body waste.
- Wash any skin that has been exposed to body waste with soap and water.
- Linens or clothing that are soiled with body fluids or body waste should be washed separately from other linens and clothing. If you do not have a washer, place the soiled linens in a plastic bag until they can be washed.
- Wash hands with soap and water after touching linens or clothing that may be soiled with body fluids.

# INTRAVENOUS CANCER TREATMENT EDUCATION



## MOGAMULIZUMAB

### Intimacy, sexual activity, contraception, and fertility

This treatment may cause changes that can affect intimacy and sexuality, including desire and body image. Maintaining physical closeness and/or intimacy with loved ones can be continued during treatment. Holding hands, hugging, and kissing can be done safely. It is recommended that you talk to your care team about any restrictions or questions you may have.

Some treatments can influence the ability to have children, also known as fertility. If you're interested in preserving fertility, talk to your care team before treatment. Ask your healthcare provider to determine when it is safe to become pregnant after your treatment. Patients of reproductive ability should not become pregnant or get their partners pregnant while receiving mogamulizumab. Some of the drugs you receive may be present in semen and vaginal secretion for many days after treatment. You should use barrier devices, such as condoms, during sexual activity to limit exposure to body fluids.

- Talk to your care team about birth control. Not all options may be right for your treatment or cancer. Effective contraception could include one or more of the following: barrier methods (e.g. condoms), hormone methods (e.g. birth control pills), or surgery.
- Tell your care team if you become pregnant or plan to breastfeed.

### Additional resources

**Product website:**

**Poteligeo:** <https://poteligeo.com/>

**Prescribing information:**

**Poteligeo:** <https://poteligeo.com/downloads/full-prescribing-information.pdf>

### Additional instructions

Updated – January 7, 2024

**Important notice:** The Association of Community Cancer Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), National Community Oncology Dispensing Association, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the therapy derived from information provided by the drug manufacturer and other resources. This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this therapy and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this therapy by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. The collaborators are not making any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual receiving therapy. All decisions related to receiving this therapy should be made with the guidance and under the direction of a qualified healthcare professional.

**Permission:** Intravenous Cancer Treatment Education (IVE) sheets are provided as a free educational resource for patients with cancer in need of concise, easy-to-understand information about intravenous cancer therapy. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the OCE website for information. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holder. Please email permission requests and licensing inquiries to [Contact@NCODA.org](mailto:Contact@NCODA.org).

Copyright © 2023 by NCODA. All rights reserved.